



Sometimes emotional difficulties can lead to feelings of despair, including thoughts about hurting yourself, ending your own life or hurting others. Speak to a friend or family member Call NHS direct on111 select option 2 'in mental health crisis' Call the Samaritans 116 123 In an emergency got to A&E for assessment Visit you GP to explore your options Please find details in your handouts.

Nature of the course

- Three webinar sessions
- Starting 6pm, aim to end at 7:15pm, with time for questions
- Evidence-based treatment programme for symptoms of panic and panic attacks
- Home tasks to support you to manage your panic symptoms
- A specific treatment programme but not 'group therapy'


4


5

Group Agreements



- Awareness and mindfulness of others.
- Focus and time.
- Confidentiality.
- No such thing as a stupid question.
- Respect each other and avoid bad language. No -isms; No insults.
- If you arrive late, just join and listen.








Symptoms of Anxiety in the Mind



- Feeling tense, nervous or unable to relax.
- Having a sense of dread, or fearing the worst.
- The world seems like it is speeding up or slowing down around you.
- Thinking others can see you are anxious, or are looking at you.
- Worrying about the future.

10



11


What is a Panic Attack?



- A sudden rush of anxiety which may exaggerate your usual anxiety symptoms.
- You may notice additional symptoms that you don't usually experience when anxious.
- Generally lasting for around 5-20 minutes.
- Can be a response to something we fear, general stress or a specific threat
- Can come 'out of the blue' when not anxious, or even asleep.








Catastrophic Misinterpretations

inclusion

- The physical symptoms of panic can be very scary when we don't know what is causing them.
- When we notice these symptoms, we try to explain them.
- Usually, we associate physical symptoms with physical health problems and these can seem dangerous.
- When we assume something dangerous is happening in the body (e.g. a heart attack), we become more anxious, keeping the panic going.

16




17

Safety Behaviours



- Because panicking can be scary, you may take steps to avoid panicking in future, or to make anxious situations more comfortable for you.
 - $\circ \quad \text{Obvious Avoidance: Completely avoiding a place, object or activity} \\ \text{that triggers symptoms of anxiety, } \textit{e.g.} \text{ avoiding spiders.}$
 - Subtle Avoidance: Not engaging in activities that may mimic symptoms
 of anxiety, e.g. exercise, or introducing new things to make you more
 comfortable, e.g. always being with others
- These behaviours provide relief from some panic symptoms, but only in the short-term.







As we discussed earlier, experiences of panic are very personal and differ from person to person. Next week will allow you to understand what is really going on in your body when you're panicking. Before you do that, it is helpful to have a short summary of the symptoms you experience. Spend some time before next session writing how you experience panic – the physical and cognitive symptoms and the safety behaviours you use to manage them.

